




CLINICAL GUIDELINE	
Resuscitation: Newborn	
Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

This document should be read in conjunction with this [DISCLAIMER](#)

Neonatal Resuscitation Algorithm – 2015 Update



© 2015 American Heart Association

WA Guidelines for newborn resuscitation

Adrenaline 1: 10 000 via ETT (1st dose; subsequent doses preferably by UVC)



- 1mL for term infant
- 0.5mL for preterm infant < 2kg or < 34 weeks gestation

If blender available commence oxygen for resuscitation at 30% and adjust as required.

Neopuff settings

- Term infant – 30 / 5 cmH₂O
- Preterm infant – 25 / 5 cmH₂O (< 2kg or < 34 weeks gestation)

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	August 2009	Last Reviewed:	1 st October 2018
Amendment Dates:		Next Review Date:	1 st October 2021
Approved by:	Neonatal Coordinating Group	Date:	23 rd October 2018
Endorsed by:	Neonatal Coordinating Group	Date:	
Standards Applicable:	NSQHS Standards:  		
Printed or personally saved electronic copies of this document are considered uncontrolled			