



CLINICAL GUIDELINE	
Venous and Arterial Access	
Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

This document should be read in conjunction with this [DISCLAIMER](#)

Venous Access

Venous access is desirable for all transported babies, and essential for ventilated babies.

- Peripheral IV's will suffice for more stable babies.
- UVC - The umbilical vein can usually be easily cannulated during the 1st week of life, and is preferred in:
 - Shocked infants, in whom peripheral IV insertion is difficult.
 - Any unstable baby.
 - Neonates requiring high glucose concentration of fluids (>12.5%).
 - Neonates requiring multiple infusions, especially inotropes, calcium, bicarbonate.
 - Double lumen UVC's are preferred.
- Intra-osseous access:
 - Rarely necessary, as umbilical venous access usually obtainable. Use in emergency situations.
 - Can administer fluids and medications at same dose as given through IV.



Arterial Access

Arterial access (UAC or peripheral arterial line) is usually indicated only in more unstable babies where blood pressure monitoring &/or frequent blood-gas sampling is desirable e.g.:

- Shocked infants.
- Extremely preterm infant (<28 weeks).
- HIE where therapeutic cooling is employed.

Note: Do not waste time (especially on country retrievals) inserting an arterial line when not essential.

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	August 2009	Last Reviewed:	1 st July 2017
Amendment Dates:		Next Review Date:	1 st July 2020
Approved by:	Neonatal Coordinating Group	Date:	26 th September 2017
Endorsed by:	Neonatal Coordinating Group	Date:	
Standards Applicable:	NSQHS Standards:  		
Printed or personally saved electronic copies of this document are considered uncontrolled			