



CLINICAL GUIDELINE	
Blood Sampling: Capillary, Venepuncture, Peripheral Arterial, UAC, UVC and CVC	
Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

This document should be read in conjunction with this [DISCLAIMER](#)

Contents

Capillary Blood Sampling	1
Venepuncture Sampling.....	2
Peripheral Arterial Catheter Sampling	3
Umbilical Catheter (UAC/UVC) Sampling.....	4
Central Venous Catheter (CVC) Sampling	4

Capillary Blood Sampling

In the absence of arterial access, capillary blood samples can be used where the total sample volume is < 1mL:


- Blood gas analysis
- U&E's/FBC/antibiotic assays/caffeine levels/SBR/PGL
- Newborn screening.
- Blood glucose monitoring
- Bone bloods

Equipment

- 1% Chlorhexidine/20% alcohol swab
- Neonatal approved lancet
- Gauze swabs/cotton wool balls
- Specimen tubes/capillary blood tube

Procedure

1. Consider pain relief such as sucrose, skin to skin holding by parents or breastfeeding.
2. Check pathology form with infant's identification bands.
3. Check that investigations are correct for that infant.
4. Collect appropriate sample tubes and equipment for blood sampling.
5. Wash hands and wear gloves.
6. Identify the puncture site that you are going to use, as per [Fig.1](#).



- Clean site and allow to dry for 30 seconds.
- Lance the skin and wipe away the first drop using a gauze swab.
- Hold the puncture site downward applying a gentle pumping action and allow blood to flow into the specimen container.
- Wipe the heel clean with a cotton ball and gently compress the site until bleeding has stopped.
- Label and check with another person the collected samples are correctly labelled at the bedside.
- Blood sampling site, the test ordered and the volume of blood taken should be documented on the patient observation chart.

Venepuncture Sampling

Venepuncture in neonates can be performed by nursing staff deemed competent in the procedure after completion of the learning package.

Venepuncture can be used for blood samples where the total sample volume is > 1mL and there is no arterial access, or where capillary sampling is not advisable. Sick neonates may require frequent intravenous cannulation it is therefore important that when performing a venepuncture the most distal aspect of the hand or the foot is used.

Equipment

- 1% Chlorhexidine/20% alcohol swab
- Venepuncture needle
- Specimen bottles
- Cotton balls

Procedure

1. Consider sucrose.
2. Check that the investigations are correct for that infant.
3. Choose area to be punctured.
4. Clean skin as per protocol.
5. Puncture skin with the bevel uppermost.
6. Direct the needle into the vein at a 45° angle.
7. Await blood return and then allow blood to drip into specimen bottle.
8. Gently remove the needle.

Blood Sampling: Capillary, Venepuncture, Peripheral Arterial, UAC, UVC and CVC

9. Maintain pressure on the site until bleeding has stopped.
10. Document sample method, the test ordered and the volume of blood taken on the observation chart.

Peripheral Arterial Catheter Sampling

Taking a peripheral arterial blood sample is a standard aseptic procedure and can be performed by staff deemed competent in this procedure.

Key Points

- Blood samples are only to be sent to the laboratory (known as a FORMAL sample) as requested by medical staff. Most blood samples can be analysed within the nursery setting (known as an INFORMAL sample).
- When analysing an informal biochemistry sample, the volume of the specimen is important. To obtain an accurate result, the sample must be 0.1mL. A smaller sample of 0.05mL is needed for a blood gas analysis.
- **Flush with 0.9% Sodium Chloride only.**

Equipment

- Blue tray
- 2 mL syringes (x2 or 3)
- Blood gas syringe
- 2% Chlorhexidine/20% alcohol swab
- 0.9% Sodium Chloride (for flush)
- Red combi stop
- Specimen bottles
- Labels
- Request form

Procedure

1. Perform hand hygiene.
2. Clean blue tray and gather equipment.
3. Perform hand hygiene.
4. Prepare equipment. Open all syringe packets, leaving the syringes lying within the packets to keep clean.
5. Prime one of the 2 mL syringes with 0.9% NaCl.
6. Place infusion pump on hold. Temporarily suspend arterial line monitoring.
7. Perform hand hygiene.
8. Don gloves.
9. Ensure 3-way tap is turned off halfway between ports. Remove and discard red combi-stop and wipe port with chlorhexidine/ alcohol swab and allow to dry. Connect empty 2 mL syringe. Open 3-way tap and gently withdraw 1 mL of blood. Turn 3-way tap off halfway between ports (This blood is to be returned to the infant post sampling).
10. Withdraw 0.05-0.1mL of blood with the heparinised blood gas syringe. Observe the infant's vital signs and TcPO₂, TcPCO₂. Turn the 3-way tap off midway, remove the gas syringe and immediately apply the filter cap. Gently expel the air from the gas syringe. Do not flick the syringe as this causes air bubbles to mix with the sample.

Blood Sampling: Capillary, Venepuncture, Peripheral Arterial, UAC, UVC and CVC

Mix sample well by inverting 4 times and rolling for 20 seconds. Analyse sample within 10 minutes. Always remix prior to analysis if a delay has occurred.

11. Take further blood samples as required.
12. Return initial blood removed back to the infant. Observe infant's limb for any signs of blanching.
13. Flush catheter with enough solution to clear most of blood from the line. This takes approximately 0.5mL.
14. Place new red combi-stop in situ. Recommence infusion pumps connected to umbilical arterial catheter. Reactivate alarms on monitor.

Umbilical Catheters (UAC/UVC) Sampling

Taking an umbilical blood sample is a standard aseptic technique and can be performed by staff deemed competent in this procedure.

Refer to [Umbilical Catheters Insertion and Management](#) for sampling procedure.

Central Venous Catheter (CVC) Sampling

Obtaining blood samples from a central venous line is a Standard Aseptic Technique which is to be performed by staff deemed competent in the procedure.

Key Points

- Sepsis is one of the most common complications of accessing CVC's. Blood sampling from CVC's should be restricted to critically ill infants who have no arterial access, as frequent blood sampling will shorten the life of the central venous catheter. **Authorization** by medical staff for sampling is to be documented in the infant's notes.
- The decision to take a blood sample from a catheter infusing medications, especially opiates or inotropes must be discussed with medical staff to avoid bolus injections of medications.
- 28G Premicath should **NOT** be used for sampling.

Equipment

- Dressing pack and 2% Chlorhexidine/ Alcohol swab
- 2 mL syringe x 1
- 10 mL syringe x 2
- Blood gas syringe, specimen bottles (if required)
- Sterile heparinised sodium chloride or heparinised glucose solution - 2 mL

Procedure


1. All solution being infused into a central venous catheter is to be heparinized.
2. Place central venous line infusion on hold.
3. Thoroughly wipe the needleless bung with 2% Chlorhexidine swab and allow to dry. Place drape from dressing pack under cleaned bung.
4. Using empty 10 mL syringe, withdraw 1 mL of fluid from the CVC to remove infusiate from the catheter.
5. Place this syringe on the dressing pack to return to infant after sample is taken.
6. Using 2 mL syringe, withdraw the required blood sample. Take a blood gas first if required. Request assistance from another staff member to perform blood gas analysis.
7. Return the blood from the 10 mL syringe to the infant.

Blood Sampling: Capillary, Venepuncture, Peripheral Arterial, UAC, UVC and CVC

8. NEVER RETURN BLOOD IF A POTASSIUM INFUSION IS IN PROGRESS.
9. Flush the catheter with 0.5-1 mL of heparinised solution in the 10 mL syringe.
10. Recommence central venous line infusion.
11. Transfer blood sample to specimen bottles. Label and verify infant's identification details.

Related CAHS internal policies, procedures and guidelines
CAHS: Infection Prevention and Control <ul style="list-style-type: none"> • Aseptic Technique

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	August 2006	Last Reviewed:	16th August 2019
Amendment Dates:		Next Review Date:	16th August 2019
Approved by:	Neonatal Coordinating Group	Date:	24th September 2019
Endorsed by:	Neonatal Coordinating Group	Date:	24th September 2019
Standards Applicable:	NSQHS Standards: 		
Printed or personally saved electronic copies of this document are considered uncontrolled			