



CLINICAL GUIDELINE

Post-Operative Care

Scope (Staff):	Nursing and Medical Staff
Scope (Area):	NICU KEMH, NICU PCH, NETS WA

This document should be read in conjunction with this [DISCLAIMER](#)

Collaborative management between surgical, medical and nursing staff is vital to achieve a positive surgical outcome.

Key Points

- Assess the infant's pain responses and ensure adequate analgesia is administered.
- The infant's vital signs should be monitored closely.
- A blood gas analysis is done on return to the unit and imbalances corrected.
- A strict fluid balance should be maintained, ensuring balance from theatre is accounted for.
- Observe the wound site(s), dressings and any drains.
- Infants' who require transportation to another hospital less than 24 hours post-surgical procedure:
 - Must be transported by the NETS team in the NETS transport cot. Nurse only is permissible if authorized by SR or Consultant.
 - Must be continuously cardiopulmonary monitored.
 - Require suction, oxygen and an appropriate sized bag and mask available **at all times** during transportation.

Minor Procedure - e.g. Hernia Repair, Eye Laser

1. Infant should receive full cardiorespiratory monitoring for 24 hours. Hourly observations for a minimum of 4 hours. Then 4 hourly for 24 hours if stable.
2. Assess the infant's pain responses and ensure adequate analgesia is administered. Hourly observations for a minimum 24 hours.
3. Thereafter the frequency of monitoring should be based on assessment of the infant's condition.

Major Procedure - Abdominal, Cardiac

1. Infant should receive full cardiorespiratory monitoring. Hourly observations for 24 hours. Hourly observations however should continue while the infant is ventilated or sedated. If ventilated consider end tidal CO₂ monitoring. Continue hourly observations if on narcotic infusion.
2. A full iSoBAR handover of the infant is given by theatre staff to the ward staff including wound sites, dressings and drains. The operation sheet MR841 should be

read by the ward staff and any specific surgical instructions noted and clarified before the theatre staff leave the unit.

3. The parents should be notified of the infant's arrival back from theatre as soon as is practical.
4. If the infant was intubated in the operating theatre, it is important to obtain a chest X-ray on return to the unit to confirm the correct placement of the endotracheal tube. A blood gas should also be done as soon as possible.
5. Surgical Drains should be managed as per surgeon's instructions. Do not attach to suction unless specifically ordered by surgeon.
6. The preterm and ex-preterm are at an increased risk of apnoea following a general anaesthetic, therefore consider caffeine therapy BUT check it has not already been administered in theatre.


Documentation

The procedure and post-operative orders should be documented on the operation sheet (MR841) by the surgeon. Any intraoperative or postoperative blood loss should be documented on the anaesthetic record MR846.2. Any drugs given in theatre should be documented on the anaesthetic record MR846.2 and the paediatric medication chart MR860.

References and related external legislation, policies, and guidelines

1. Boxwell G. Neonatal Intensive Care Nursing. 2nd ed. London: Routledge; 2010
2. Hansen A, Puder M. Manual of Neonatal Surgical Intensive Care, 2nd Edition, 2009. People's Medical Publishing House, Shelton, Connecticut.

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Neonatology		
Reviewer / Team:	Neonatal Coordinating Group		
Date First Issued:	July 2006	Last Reviewed:	1 st January 2015
Amendment Dates:		Next Review Date:	1 st January 2018
Approved by:	Neonatal Coordinating Group	Date:	January 2015
Endorsed by:	Neonatal Coordinating Group	Date:	January 2015
Standards Applicable:	NSQHS Standards: 		
Printed or personally saved electronic copies of this document are considered uncontrolled			