


Bronchiolitis

What is it?

Bronchiolitis is a flu-like illness that affects infants in the first year of life. It is caused by one of several viruses, the most common one being RSV (respiratory syncytial virus), which affects the small airways (bronchioles) in the lungs. Bronchiolitis occurs frequently in the winter months and rarely in summer. This condition starts as a simple cold and progresses to a more complicated respiratory illness with symptoms such as cough, wheeze, fast breathing and feeding difficulties.

Diagnosis

Nasal secretions may be sent to the laboratory to confirm what type of virus your child has. However, there is no special test to confirm the diagnosis.



Home Management

Always try not to smoke in the home or around your baby. This is especially important around babies with any respiratory illness.

- Medicines do not usually help babies with bronchiolitis. Antibiotics are not given because bronchiolitis is caused by a virus and antibiotics do not cure viruses.
- Encourage rest.
- Give shorter breast feeds or formula, and water more frequently. This way your child does not get too tired when feeding. If your child does not get enough fluids they can get dehydrated.
- You can give Paracetamol (eg. Panadol, Dymadon, Panamax) if your child is irritable.
- Avoid contact with other babies in the first few days, as bronchiolitis is an infectious disease.

If your baby is distressed and having trouble feeding, they may need to be admitted to hospital. Staff may need to:

- Watch them closely to ensure they do not become more unwell.
- Give them some extra oxygen.
- Give them extra fluids through a drip into a vein or tube feed through the nose into the stomach.



Follow-up

Make an appointment for your child to see a doctor (or return to the Emergency Department) if they have any of the following:

- Difficulty breathing (very fast or not regular breaths).
- Turns blue or has skin that is pale and sweaty.
- A cough that is getting worse.
- Taking less than half their normal feeds or are refusing food or fluids.
- Seem very tired or are more sleepy than usual.
- If you are worried in any way.

Key points to remember

- Babies need to rest and drink small amounts more often.
- Bronchiolitis is an infectious disease in the first few days of illness.
- It is more common in babies under six months old.
- Babies are usually sick for three to five days and then recover over the next seven to 10 days. The cough may continue for two to four weeks.
- Smoking in the home increases the chance of babies having bronchiolitis and makes it worse.



Government of **Western Australia**
Child and Adolescent Health Service

This document can be made available in alternative formats on request for a person with a disability.

Child and Adolescent Health Service
15 Hospital Avenue, Nedlands, WA, 6009
Telephone: (08) 6456 2222
Produced by Emergency Department
Ref: 529 © CAHS 2016

Disclaimer: This publication is for general education and information purposes. Contact a qualified healthcare professional for any medical advice needed.
© State of Western Australia, Child and Adolescent Health Service