

Immune Thrombocytopenic Purpura (ITP)

What is ITP?

ITP is a bleeding disorder where there are fewer platelets in the blood than usual. Platelets help to clot the blood. If a child has a shortage of platelets they will bruise easily and may have other bleeding problems.

ITP occurs in around one out of every 10,000 children. The risk of any serious complication from ITP is rare.

In most children the condition will settle down quickly. The platelet count usually returns to safe levels within a few weeks. In a small number of children, the platelet count does not return to normal, even after 6 to 12 months, and further investigation may be required.

What causes ITP?

The cause is unknown, but it can follow a viral infection.

Signs and symptoms

- Bruising
- Nosebleeds or mouth bleeding
- Red pinpoint spots on the skin (petechial rash)

Diagnosis

ITP is usually diagnosed with a blood test called a 'full blood count'. The sample of blood is examined under a microscope to rule out other conditions that may cause similar symptoms to ITP.

Management

Most children will be able to be discharged home with no treatment; however your doctor will discuss the best management option for your child. The condition will get better spontaneously and your child can be monitored by your GP in conjunction with PCH.

You will need to attend the hospital and your GP to monitor the progress of the condition. It is important that you come to all follow up appointments.

Treatment is usually for patients who have serious bleeding.

- Steroids raise the platelet count quickly for a short period of time.



- Intravenous gamma globulin may be given when bleeding is life threatening

Care at home

- Children with ITP should not take aspirin or ibuprofen (eg Nurofen[®]), as this may increase bleeding. Paracetamol (e.g. Panadol[®]) is safe.
- While your child has ITP, they should attend school but they should not do activities which might cause bruising or bleeding (e.g. playing on climbing equipment, trampolines, skateboards, cycling, contact sports, and rough physical activity).

When to seek help

When your child is sent home you will be given a clinic appointment for review at the hospital within two weeks, and an emergency number.

If your child has a minor injury such as a graze to a knee or a small cut you should use normal first aid measures such as RICE (**R**est, **I**ce, **C**ompression, **E**levation).

You should return to hospital in the following circumstances:

- A prolonged (over 15 minutes) nosebleed which will not stop despite pinching the nose
- Prolonged bleeding from the mouth, gums, cheeks or throat
- Coughing up blood
- Painful muscles or joints
- Following a heavy blow to the head
- Persistent or severe headache
- Vomiting or drowsiness
- If you think your child is seriously unwell

Further information is available from the following resource:

[UK ITP Support - http://www.itpsupport.org.uk/childhooditp.htm](http://www.itpsupport.org.uk/childhooditp.htm)



Government of **Western Australia**
Child and Adolescent Health Service

This document can be made available in alternative formats on request for a person with a disability.

Child and Adolescent Health Service
15 Hospital Avenue, Nedlands, WA, 6009
Telephone: (08) 6456 2222
Produced by Emergency Department
Ref: 173 © CAHS 2017

Disclaimer: This publication is for general education and information purposes. Contact a qualified healthcare professional for any medical advice needed.
© State of Western Australia, Child and Adolescent Health Service