

Adenoidectomy

Discharge advice

Bleeding

Observe for any fresh bleeding (bright red blood) from the nose or mouth. Seek medical advice immediately if bleeding occurs.

Due to the risk of bleeding:

- a child **under 12 months** is not to travel out of the metropolitan area for **48 hours** after the operation
- a child **over 12 months** is not to travel out of the metropolitan area for **24 hours** after the operation.

Pain management

- Give either paracetamol (Panadol) or ibuprofen (Nurofen) for pain if required, as per manufacturer's instructions.
- Next dose of paracetamol can be given at _____.
- Next dose of ibuprofen can be given at _____.
- Do not give any medicine with paracetamol in it more than four times in 24 hours.
- Do not give any medicine with ibuprofen in it (e.g. Nurofen) more than three times in 24 hours.
- Do not give Aspirin, Disprin or other products containing aspirin.

Care at home

- Do not blow the nose for seven days, wipe gently instead.
- Your child may experience bad breath. This is normal and should be gone within 10 to 14 days. Use mouth wash to manage bad breath.
- If your child develops a fever please see your family doctor.
- Your child can resume a normal diet and fluids.

Daily activities

- Your child should not play sport or attend school for seven days after their surgery.
- Your child should not go swimming due to the risk of infection for two weeks after their surgery.



Advice following an anaesthetic

Food and Drink

- When your child is tolerating clear fluids, gradually increase the amount they are being given.
- When your child is hungry, a light diet is recommended. Avoid greasy or fatty foods.
- A normal diet can be resumed the day after a procedure.

Vomiting

- If your child vomits, give nothing for half an hour, then start sips of clear fluids. Gradually increase the amount being given when your child is tolerating fluids.
- Seek medical advice if vomiting continues.

Directions

- Seek medical advice if your child has problems passing urine.
- A responsible adult must stay with the child for at least the first 24 hours after a procedure.
- Seek medical advice if the area where your child's intravenous catheter was put in has increasing redness, swelling or pain.

Contact details

If you have any concerns after your child's discharge from hospital, please phone 6456 2222 and ask for the Surgical Short Stay Unit between 7am and 9pm, Monday to Friday. After these hours ask for Ward 4B or contact your GP.

To cancel or change your appointment, or if you've forgotten your appointment details, please call Outpatient Direct on 1300 855 275.



Government of **Western Australia**
Child and Adolescent Health Service

15 Hospital Avenue, Nedlands, WA, 6009
Telephone: (08) 6456 2222
Produced by Surgical Short Stay Unit
Ref: 118 © CAHS 2017 rev 2020



Disclaimer: This publication is for general education and information purposes. Contact a qualified healthcare professional for any medical advice needed.
© State of Western Australia, Child and Adolescent Health Service

This document can be made available in alternative formats on request for a person with a disability.