



## GUIDELINE

# Apnoea of Prematurity

<b>Scope (Staff):</b>	Nursing and Medical Staff
<b>Scope (Area):</b>	NICU KEMH, NICU PCH, NETS WA

### Child Safe Organisation Statement of Commitment

CAHS commits to being a child safe organisation by applying the National Principles for Child Safe Organisations. This is a commitment to a strong culture supported by robust policies and procedures to reduce the likelihood of harm to children and young people.

This document should be read in conjunction with this [disclaimer](#)

## Aim

To provide Neonatology staff guidance in the management and identification of infants at risk of apnoea of prematurity.

## Risk

Delayed management and increasing the risk of premature infants to having apnoea and hypoxic episodes.

## Definition

Absence of breathing in the premature infant for a period of > 15 seconds often associated with bradycardia and/or desaturation.

## Epidemiology

The incidence of apnoea of prematurity is inversely correlated with gestational age. Recurrent apnoea and bradycardia occurs in most infants under 30 weeks gestation, in about 50% of infants at 30-32 weeks gestation, and only about 10% of those at 34-36 weeks gestation.

Apnoea of prematurity is primarily due to immaturity of the respiratory centre and control of the upper airway; most cases are due to a combination of central and obstructive apnoea.

It is important to manage the following additional factors which can either cause or accentuate apnoea:

- Systemic illness - sepsis, hypoxemia, shock.
- CNS – IVH, seizures, asphyxia, cerebral malformations.

- CVS conditions: e.g. PDA
- Anaemia
- Drugs - opiates, sedation.
- Post-general anaesthesia.
- Metabolic disturbances: hypoglycaemia, hypocalcaemia, inborn error of metabolism.
- Hypothermia, hyperthermia.
- Obstructive respiratory problems laryngo-tracheomalacia, macroglossia, micrognathia, , choanal atresia
- Gastro-oesophageal reflux.

## Management

- Address any underlying factor that may be aggravating apnoeic episodes.
- Monitor infants with cardio-respiratory monitoring and oxygen saturation. Refer to [Monitoring and Observation Frequency](#).
- If an episode of apnoea/bradycardia does not resolve spontaneously, gentle tactile stimulation maybe required.
  - If response is slow or cyanosis present, then bag and mask ventilation may be needed.
- Prone positioning stabilises the chest wall, reducing the frequency of apnoea.
- CPAP should be optimised. Occasionally non-invasive ventilation (NIPPV) may be required. Intubation is sometimes required when the above fails.
- **Caffeine** is a potent respiratory and CNS stimulant and is the pharmacological agent of choice. It has been shown to decrease apnoea and reduce the need for ventilator support. It also reduces short term morbidities such as BPD and ROP. It is cost-effective and improves long term outcomes with lower rates of developmental co-ordination disorder.
  - For dosage and side effects refer to Neonatal Medication Protocols - [Caffeine](#).

## Who to Treat

- Infants < 30 weeks gestation are generally prescribed caffeine routinely. Caffeine has been shown to reduce post-extubation apnoea.
- Infants 30-34 weeks gestation are generally prescribed caffeine if they are symptomatic.

- Consider prescribing caffeine for a neonate in the immediate post-operative period following a general anaesthetic, or prior to immunisations in an ex-preterm neonate.



## Stopping Caffeine

There are no trial data to support decisions to cease treatment. Apnoea of prematurity decreases with advancing age and can generally be expected to resolve by 37 weeks corrected gestational age (CGA). Apnoea which continues beyond this period is often multifactorial (see above).

- Caffeine can generally be ceased between 32- and 34-weeks' gestation, provided the baby has been stable. Some extremely preterm neonates will require treatment for longer.
- Once the infant has ceased caffeine, infants should remain monitored with an oximeter for a minimum of 5 days and for 48 hours after the last apnoea/bradycardia. Refer to [Monitoring and Observation Frequency](#).

In general, we do not measure caffeine levels routinely.

This document can be made available in alternative formats on request.

Document Owner:	Neonatology		
Reviewer / Team:	Neonatology Coordinating Group		
Date First Issued:	October 2006	Last Reviewed:	August 2022
Amendment Dates:		Next Review Date:	August 2025
Approved by:	Neonatology Coordinating Group	Date:	23 <sup>rd</sup> August 2022
Endorsed by:	Neonatology Coordinating Group	Date:	
Standards Applicable:	NSQHS Standards:   Child Safe Standards: 1,10		

Printed or personally saved electronic copies of this document are considered uncontrolled



## Healthy kids, healthy communities

Compassion
Excellence
Collaboration
Accountability
Equity
Respect

Neonatology | Community Health | Mental Health | Perth Children's Hospital